MycoKeys 111: 1–20 (2024)
DOI: 10.3897/mycokeys.111.137347

Research Article

Taxonomic study of *Hydnoporia* (Hymenochaetales, Hymenochaetaceae) in East Asia with two new species

Minseo Cho[1*], Yoonhee Cho[2*], Sun Lul Kwon[1,3], Dohye Kim[2], Kentaro Hosaka[4], Young Woon Lim[2], Jae-Jin Kim[1]

1 *Division of Environmental Science and Ecological Engineering, College of Life Sciences and Biotechnology, Korea University, 145 Anam-ro, Seongbuk-gu, Seoul, Republic of Korea*
2 *School of Biological Sciences and Institute of Biodiversity, Seoul National University, Gwanak-ro, Gwanak-gu, Seoul, Republic of Korea*
3 *BK21 FOUR R&E Center for Environmental Science and Ecological Engineering, Korea University, 145 Anam-ro, Seongbuk-gu, Seoul, Republic of Korea*
4 *Department of Botany, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan*

Corresponding authors: Young Woon Lim (ywlim@snu.ac.kr); Jae-Jin Kim (jae-jinkim@korea.ac.kr)

Abstract

The genus *Hydnoporia* (Hymenochaetales, Hymenochaetaceae) was first described by Murrill in 1907. However, species of *Hydnoporia* were subsequently reclassified into the genera *Hymenochaete*, *Hymenochaetopsis*, and *Pseudochaete* due to overlapping morphological characteristics, leading to multiple synonyms and confusion among researchers. Recent phylogenetic analyses based on multimarker datasets have clarified the diversity and relationships within *Hydnoporia*, but East Asian species remain underrepresented due to limited morphological data. To address this gap, we conducted a comprehensive morphological and phylogenetic analysis of East Asian *Hydnoporia* specimens using internal transcribed spacer (ITS) and translation elongation factor 1–α (*tef1*) regions. From 42 specimens, we identified six species, including two novel species, *Hydnoporia orienticorrugata* **sp. nov.** and *Hydnoporia subtabacina* **sp. nov.**, and we report *Hydnoporia rimosa* for the first time in Korea. Phylogenetic analyses also support the recombination of *Hymenochaete intricata* and *Hymenochaetopsis rigidula* as *Hydnoporia intricata* **comb. nov.** and *Hydnoporia rigidula* **comb. nov.**, respectively. By elucidating the phylogenetic relationships and morphological traits of *Hydnoporia* species from East Asia, this study contributes to a deeper understanding of the global diversity and phylogeny of the genus.

Key words: *Hymenochaete*, *Hymenochaetopsis*, novel species, phylogeny, *Pseudochaete*

OPEN ACCESS

Academic editor: R. Henrik Nilsson
Received: 18 September 2024
Accepted: 1 November 2024
Published: 25 November 2024

Citation: Cho M, Cho Y, Kwon SL, Kim D, Hosaka K, Lim YW, Kim J-J (2024) Taxonomic study of *Hydnoporia* (Hymenochaetales, Hymenochaetaceae) in East Asia with two new species. MycoKeys 111: 1–20. https://doi.org/10.3897/mycokeys.111.137347

Copyright: © Minseo Cho et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

The genus *Hydnoporia* Murill. in the family Hymenochaetaceae (Hymenochaetales, *Basidiomycota*) was first described in 1907 and typified through *Hyd. olivacea* (Schwein.) Teixeira (≡*Sistotrema fuscescens*). *Hydnoporia* is characterized by brownish basidiomes of diverse shapes, ranging from effused-reflexed to pileate, cylindrical to allantoid basidiospores, and setae of various sizes (Miettinen et al. 2019). *Hydnoporia* species are found across Asia, Europe, and

* These authors contributed equally to this work.

North America (He and Li 2013; Miettinen et al. 2019; Palla et al. 2023). They inhabit the branches or stumps of conifer and deciduous trees and are classified as wood-decaying corticioid fungi (Miettinen et al. 2019). Until the 1990s, many corticioid species had been delimited based on their morphological traits (Parmasto 1995; Léger 1998). Due to their overlapping morphological characteristics, *Hydnoporia* species were previously placed under the genus *Hymenochaete* Lev., which includes species characterized by brown-colored basidiomes, small cylindrical basidiospores, and the presence of setae (Parmasto et al. 2014; Miettinen et al. 2019). The lack of distinguishable characteristics between the two genera prevented a clear separation despite the efforts of some taxonomists (Parmasto 1995; Léger 1998).

Technological advancements made DNA-based analyses possible during the 1990s and onwards. This led to many taxonomic revisions, including the division of *Hymenochaete* into two separate clades: *Hymenochaete sensu stricto* and *Pseudochaete* T. Wagner & M. Fisch (Wagner and Fischer 2002). However, the genus *Pseudochaete* was present under the botanical code in the green algae *Pseudochaete* W. West & G. S. West 1903, which was considered repetitious for microorganisms despite differences in lineages. For this reason, a new name *Hymenochaetopsis* S. H. He & Jiao Yang was introduced (Yang et al. 2016). However, none of these studies considered *Hydnoporia*, which had priority. *Hymenochaetopsis* and *Pseudochaete* eventually became synonyms of *Hydnoporia* (Miettinen et al. 2019). Additionally, the clade of *Hymenochaete tabacina* (Sowerby) Lèv. and its neighboring species were encompassed within *Hydnoporia* as it formed a sister clade to the type species, *Hyd. olivacea* (Miettinen et al. 2019; Wu et al. 2022).

Currently, 14 species are accepted in *Hydnoporia*, and 20–27 more species candidates have been revealed through multimarker-based phylogenetic analysis (Miettinen et al. 2019). However, previous studies did not fully cover East Asian specimens owing to the lack of detailed morphological data, and this was problematic because East Asia is expected to be the most diverse region for *Hydnoporia* species (He and Li 2013; Miettinen et al. 2019; Palla et al. 2023). To fully understand the global diversity of *Hydnoporia*, a detailed morphological and phylogenetic analysis of East Asian specimens is essential. This study assesses *Hydnoporia* specimens from the Republic of Korea and Japan based on multigenetic marker phylogeny (ITS+*tef1*) and morphological characteristics, and it describes two new species, *Hyd. orienticorrugata* sp. nov. and *Hyd. subtabacina* sp. nov. It furthermore notes one unrecorded species from the Republic of Korea, *Hyd. rimosa*. Additionally, we propose synonymizing *Hymenochaete intricata* and *Hymenochaetopsis rigidula* as *Hyd. intricata* and *Hyd. rigidula*, respectively, considering the phylogenetic relationship of these species with *Hydnoporia*.

Materials and methods

Specimen collection

We studied 43 specimens labeled as *Hydnoporia*, *Hymenochaete*, *Hymenochaetopsis*, and *Pseudochaete* deposited in the Korea University Culture Collection (**KUC**), Seoul National University Fungus Collection (**SFC**), and National Institute of Biological Resources (**NIBR**) in this study. Specimens were collected nationwide from 2012 to 2021 and stored with silica gel under dry conditions to

prevent contamination. A *Hyd. yasudai* specimen from **TNS** (Mycological Herbarium of the Department of Botany, National Museum of Nature and Science, Tsukuba, Japan) was also included.

Molecular identification

Genomic DNA was extracted from the dried specimens using AccuPrep® Genomic DNA Extraction Kit (Bioneer, Daejeon, Korea). Primer sets ITS1F/ITS4, ITS1F/ITS4B, ITS5/ITS4, or ITS5/LR3 were used to amplify the nuclear ribosomal internal transcribed spacer (ITS) region (White et al. 1990; Gardes and Bruns 1993). For the translation elongation factor 1−α (*tef1*) region, the primer set 983F/1567R was used (Rehner and Buckley 2005). PCR products were purified using the AccuPrep® PCR Purification Kit following the manufacturer's instructions. DNA sequencing was performed by Cosmogenetech (Seoul, Korea) using the amplified products. All sequencing results were edited using the SeqMan Lasergene package version 7.0.0 (DNAStar Inc., Madison, WI). The newly generated sequences were deposited in GenBank (Sayers et al. 2024; Table 1).

Reference ITS and *tef1* sequences for phylogenetic analysis were obtained from GenBank following Miettinen et al. (2019). All sequences were aligned by region using MAFFT v. 7.490 (Katoh and Standley 2013). Sequence alignment and concatenation were performed using Geneious Prime 2023.2.1. (https://www.geneious.com; Suppl. materials 1, 2). A phylogenetic tree was inferred on the CIPRES web portal using the concatenated ITS and *tef1* datasets with maximum likelihood (ML) and Bayesian inference (BI) methods (Miller et al. 2015). For tree inference, ITS and *tef1* sequences were partitioned into eight regions: ITS1, 5.8S, and ITS2 for the ITS region and exon 1, intron 1, exon 2, intron 2, and exon 3 for *tef1* region. jModeltest v. 2.1.10 was used to select the best-fitting substitution model for ITS and *tef1* regions (Darriba et al. 2012). The best-fitting models for ITS1, 5.8S, ITS2, exon 1, intron 1, exon 2, intron 2, and exon 3 regions were HKY+G, JC, HKY+G, K80+G, GTR+I, K80+I, HKY+I, and K80+G, respectively. The ML tree was inferred using RAxML-HPC2 in XSEDE v. 8.2.12 with 1,000 bootstrap replicates (Stamatakis 2014). BI analyses were conducted using MrBayes v. 3.2.3 on XSEDE, with the best model selected for each marker by sampling every 1,000 generations for 20 million generations (Ronquist and Huelsenbeck 2003). The phylogenetic tree was edited using FigTree v. 1.4.3 (Rambaut 2018) and Adobe Illustrator CS6 (Adobe Systems Inc., San Jose, CA, USA).

Morphological observation

Macroscopic images of each specimen were captured using a Sony α 6500 camera (Sony, Tokyo, Japan). Microscopic structures were observed using an Olympus BX51 light microscope (Olympus, Tokyo, Japan) at 40–1000 × magnification. The images were captured using a DP20 microscope (Olympus, Tokyo, Japan). At least 20 basidiospores, basidia, and setae were examined to measure the size. Specific color terms were obtained from the Munsell Soil Color Book (Color 2009). The following abbreviations were used: **L** = mean spore length; **W** = mean spore width; **Q** = L/W ratio; **x** = the number of basidiospores measured; **y** = the number of specimens; and **n** = x/y.

Table 1. List of analyzed *Hydnoporia* specimens with GenBank accession numbers of ITS and *tef1* sequences.

Species name	Specimen voucher	Country	GenBank accession no.[a]		References
			ITS	*tef1*	
Hydnoporia corrugata	Jon Klepsland 2011-7-24 (O F-247869)[T]	Norway	MK514613	MK552138	(Miettinen et al. 2019)
	KCG001	Ireland	JQ246338	–	(Grundy et al. 2012)
	A6_wood_6	Great Britain	JN230421	–	(Grundy et al. 2012)
	B1_wood_inner	Ireland	JN230422	–	(Grundy et al. 2012)
	A3_wood_2	Great Britain	JN230419	–	(Grundy et al. 2012)
Hyd. diffissa	Otto Miettinen 19463 (H 7008917)[T]	USA, North Carolina	MK514611	MK552136	(Miettinen et al. 2019)
	Otto Miettinen 17127.4 (H)	USA, New York	MK514598	–	(Miettinen et al. 2019)
Hyd. gigasetosa	He1461	China, Yunnan	KT828671	–	(Yang et al. 2016)
	He1442	China, Yunnan	KT828670	–	(Yang et al. 2016)
***Hyd. intricata* comb. nov.**	**KUC20121123-03**	**Korea**	**PP992254**	**PQ066850**	**This study**
	KUC20210428-20	**Korea**	**PP992255**	**PQ066851**	**This study**
	KUC20211030-01	**Korea**	**PP992256**	**PQ066852**	**This study**
	SFC20120820-11	**Korea**	**PP992257**	**–**	**This study**
	SFC20140313-01	**Korea**	**PP992258**	**PQ066853**	**This study**
	SFC20160920-36	**Korea**	**PP992259**	**PQ066854**	**This study**
	SFC20170822-68	**Korea**	**PP992260**	**PQ066855**	**This study**
	SFC20170908-28	**Korea**	**PP992261**	**PQ066856**	**This study**
	He1181	China	JQ279609	–	(He and Dai 2012)
	He412	China	JQ279608	–	(He and Dai 2012)
	He21064	China	KC505556	–	Unpublished
Hyd. lamellata	Cui7629	China	JQ279603	–	(He and Dai 2012)
	Dai10527	China	JQ279605	–	(He and Dai 2012)
Hyd. laricicola	Viacheslav Spirin 5400 (H)	Russia, Khabarovsk	MK514606	MK552132	(Miettinen et al. 2019)
	Dai13458[T]	China, Heilongjiang	KT828672	–	(Yang et al. 2016)
	Dai11046	China, Nei Mongol	JQ279616	–	(He and Dai 2012)
	Wu 1207-122	China, Jilin	KT828673	–	(Yang et al. 2016)
Hyd. latesetosa	He492	China, Hainan	JQ716404	–	(He and Li 2013)
	He502[T]	China, Hainan	NR_120093	–	(He and Li 2013)
Hyd. olivacea	Otto Miettinen & Kelo Käppi 16044 (H 7005770)[T]	USA, Massachusetts	MK514612	MK552137	(Miettinen et al. 2019)
	P1201B	Peru	EU977192	–	(Smith et al. 2008)
	CMH529	USA, Missouri	KF800618	–	(Rittenour et al. 2014)
	f2Fc06	USA, Texas	GU721341	–	(Noris et al. 2011)
	319	Antarctica	KC785573	–	(Connell and Staudigel 2013)
	CFMR:DLL2011-223	USA, Wisconsin	KJ140712	–	(Brazee et al. 2014)
	CBS:126040	USA, North Carolina	MH864055	–	(Vu et al. 2019)
Hyd. rhododendri	N. Gerhold 2005-6-3	Austria	MK514593	–	(Miettinen et al. 2019)
	Viacheslav Spirin 6476 (H)	Russia, Primorsky Krai	MK514599	MK552127	(Miettinen et al. 2019)
	Viacheslav Spirin 6450 (H)	Russia, Primorsky Krai	MK514603	–	(Miettinen et al. 2019)
***Hyd. rigidula* comb. nov.**	SFC20140314-10	Korea	KX792928	–	(Kim et al. 2016)
	SFC20140411-08	Korea	KX792929	–	(Kim et al. 2016)
	SFC20140411-20	Korea	KX792930	–	(Kim et al. 2016)
	SFC20140703-24	Korea	KX792931	–	(Kim et al. 2016)
	SFC20140723-16	**Korea**	KX792932	**PQ066857**	(Kim et al. 2016), **This study**
	SFC20160713-06	**Korea**	**PP992262**	**PQ066858**	**This study**
	SFC20170324-10	**Korea**	**PP992263**	**PQ066859**	**This study**
	He379	China	JQ279613	–	(He and Dai 2012)
	He343	China	JQ279612	–	(He and Dai 2012)

Species name	Specimen voucher	Country	GenBank accession no.[a]		References
			ITS	*tef1*	
Hyd. rimosa	**KUC20121109-19**	**Korea**	**PP992264**	**–**	**This study**
	Viacheslav Spirin 5277 (H)	Russia, Khabarovsk	MK514592	MK552122	(Miettinen et al. 2019)
	Viacheslav Spirin 5678 (H)	Russia, Khabarovsk	MK514594	MK552123	(Miettinen et al. 2019)
	Viacheslav Spirin 6104 (H)	Russia, Khabarovsk	MK514595	MK552124	(Miettinen et al. 2019)
Hyd. subrigidula	He1123	China, Yunnan	JQ716402	–	(He and Li 2013)
	He1157^{T}	China, Yunnan	NR_120092	–	(He and Li 2013)
Hyd. tabacina	A. M. Ainsworth 2017-1-17 (K(M) 233332)	Great Britain	MK514614	MK890223	(Miettinen et al. 2019)
	Otto Miettinen 22126 (H)	Finland	MK782755	MK787232	(Miettinen et al. 2019)
	Viacheslav Spirin 6066a (H)	Russia, Nizhny Novgorod	MK514600	MK552128	(Miettinen et al. 2019)
Hyd. tabacinoides	CLZhao986	China, Yunnan	MG231566	–	Unpublished
	CLZhao859	China, Yunnan	MG231565	–	Unpublished
	Cui10428	China	JQ279604	–	(He and Dai 2012)
Hyd. yasudai	**KUC20100409-18**	**Korea**	**PP992265**	**PQ066860**	**This study**
	KUC20180326-05	**Korea**	**PP992266**	**PQ066861**	**This study**
	KUC20210319-14	**Korea**	**PP992267**	**PQ066862**	**This study**
	SFC20150707-58	**Korea**	**PP992268**	**PQ066863**	**This study**
	SFC20150902-19	**Korea**	**PP992269**	**–**	**This study**
	SFC20160114-04	**Korea**	**PP992270**	**PQ066864**	**This study**
	SFC20160512-38	**Korea**	**PP992271**	**–**	**This study**
	SFC20160517-06	**Korea**	**PP992272**	**PQ066865**	**This study**
	SFC20160527-41	**Korea**	**PP992273**	**PQ066866**	**This study**
	SFC20160614-52	**Korea**	**PP992274**	**–**	**This study**
	SFC20160712-18	**Korea**	**PP992275**	**PQ066867**	**This study**
	SFC20180410-24	**Korea**	**PP992276**	**PQ066868**	**This study**
	SFC20180712-04	**Korea**	**PP992277**	**PQ066869**	**This study**
	KUC11055	Korea	KJ713999	–	(Jang et al. 2015)
	KoLRI48661	Korea, Jeju	MT586954	–	Unpublished
	KoLRI_EL005212	Korea, Jeju	MN844835	–	Unpublished
	KoLRI_EL005068	Korea, Jeju	MN844834	–	Unpublished
	TNS-F78711	**Japan**	**PP992278**	**PQ066870**	**This study**
	IFO 4969	Japan	AY558598	–	(Jeong et al. 2005)
	Viacheslav Spirin 5533 (H)	Russia, Khabarovsk	MK514597	MK552126	(Miettinen et al. 2019)
	Viacheslav Spirin 6475 (H)	Russia, Primorsky Krai	MK514609	MK552135	(Miettinen et al. 2019)
	CLZhao1495	China, Yunnan	MG231611	–	Unpublished
	CLZhao1475	China, Yunnan	MG231609	–	Unpublished
	CLZhao1486	China, Yunnan	MG231610	–	Unpublished
	CLZhao1422	China, Yunnan	MG231607	–	Unpublished
	He273	China	JQ279614	–	(He and Dai 2012)
	He375	China	JQ279615	–	(He and Dai 2012)
	CLZhao867	China, Yunnan	MG231606	–	Unpublished
	CLZhao933	China, Yunnan	MH114725	–	Unpublished
	CLZhao853	China, Yunnan	MG231605	–	Unpublished
	CLZhao1549	China, Yunnan	MG231612	–	Unpublished

Species name	Specimen voucher	Country	GenBank accession no.[a]		References
			ITS	*tef1*	
***Hyd. orienticorrugata* sp. nov.**	KUC20121019-16	Korea	KJ668528	–	(Jang et al. 2016)
	KUC20121123-05	**Korea**	**PP992279**	**–**	**This study**
	KUC20131001-21	**Korea**	**PP992280**	**PQ066871**	**This study**
	SFC20140412-06	**Korea**	**PP992281**	**–**	**This study**
	SFC20150212-01	**Korea**	**PP992282**	**PQ066872**	**This study**
	SFC20150319-12	**Korea**	**PP992283**	**–**	**This study**
	SFC20150513-06	**Korea**	**PP992284**	**–**	**This study**
	SFC20151030-12[T]	**Korea**	**PP992285**	**PQ066873**	**This study**
	SFC20190619-11	**Korea**	**PP992286**	**–**	**This study**
	CLZhao938	China, Yunnan	MH114693	–	Unpublished
	He761	China	JQ279606	–	(He and Dai 2012)
	He839	China	JQ279607	–	(He and Dai 2012)
***Hyd. subtabacina* sp. nov.**	**SFC20190322-02[T]**	**Korea**	**PP992287**	**PQ066874**	**This study**
	SFC20190510-01	**Korea**	**PP992288**	**PQ066875**	**This study**
	SFC20190619-15	**Korea**	**PP992289**	**–**	**This study**
	Heikki Kotiranta 20797 (H)	Russia, Perm	MK514591	MK552121	(Miettinen et al. 2019)
	Heikki Kotiranta 25205 (H)	Russia, Kransoyarsk	MK514596	MK552125	(Miettinen et al. 2019)
	Otto Miettinen 17028.3 (H)	USA, New York	MK514601	MK552129	(Miettinen et al. 2019)
	Viacheslav Spirin 5196 (H)	Russia, Khabarovsk	MK514602	MK552130	(Miettinen et al. 2019)
	Viacheslav Spirin 6582 (H)	Russia, Khabarovsk	MK514604	–	(Miettinen et al. 2019)
	Viacheslav Spirin 6566 (H)	Russia, Khabarovsk	MK514605	MK552131	(Miettinen et al. 2019)
	Viacheslav Spirin 6520 (H)	Russia, Khabarovsk	MK514607	MK552133	(Miettinen et al. 2019)
	Viacheslav Spirin 6507 (H)	Russia, Khabarovsk	MK514608	MK552134	(Miettinen et al. 2019)
	CFMR:DLL2011-152	USA, Wisconsin	KJ140652	–	(Brazee et al. 2014)
	CFMR:DLL2011-071	USA, Wisconsin	KJ140591	–	(Brazee et al. 2014)
	CFMR:DLL2011-175	USA, Wisconsin	KJ140671	–	(Brazee et al. 2014)
	He810	China	JQ279611	–	(He and Dai 2012)
	He390	China	JQ279610	–	(He and Dai 2012)
Porodaedalea alpicola	Cui12280	China	ON358110	ON631040	(Wu et al. 2019)

[a]The sequences generated in this study are shown in bold.
[T]Indicate the type materials.

Results

Phylogenetic analyses

Based on ITS sequence analysis, the 43 assessed specimens were phylogenetically grouped within *Hydnoporia* (Fig. 1). Phylogenetic analysis of the ITS and *tef1* regions identified these specimens as six distinct Korean species (Fig. 2, Suppl. material 3). Among them, three were confirmed as previously recorded species in Republic of Korea: *Hymenochaete intricata* (Lloyd) T. Ito, *Hym. rigidula* Berk. & M.A. Curtis, and *Hydnoporia yasudai* (Imazeki) Spirin & Miettinen. One species was identified as a new record for Korea: *Hyd. rimosa* (Lloyd) Spirin & Miettinen. These four species were also well supported by morphological characteristics.

The remaining two species, previously labelled as "*Hyd. corrugata*" and "*Hyd. tabacina*", formed distinct clades from their close relatives *Hyd. corrugata*



Figure 1. BI tree based on the ITS sequence datasets. The node numbers indicate the bootstrap support values (BS) above 70% and posterior probabilities (PP) over 0.7 as BS/PP. *Hydnoporia* specimens examined in this study are shown in bold. Newly generated sequences in this study are shown in blue and bold. *Porodaedalea alpicola* (Cui 12280) is used as an outgroup. Letter codes after specimen voucher indicate ISO 3166 country code followed by the origin province. Detailed information is in Table 1. Type specimens are indicated with "T".

and *Hyd. tabacina*, respectively. However, East Asian and American "*Hyd. tabacina*" were poorly differentiated from European *Hyd. tabacina* and *Hyd. rhododendri* in the ITS phylogeny. Morphological comparison and multimarker-based phylogenetic inference with other *Hydnoporia* species support the recognition of these two as new species, which we propose as *Hyd. orienticorrugata* sp. nov. and *Hyd. subtabacina* sp. nov. Morphological descriptions of the new species are provided in the Taxonomy section.

Additionally, two species previously classified as *Hymenochaete* formed strongly supported clades within *Hydnoporia*. Therefore, we propose their reclassification as *Hyd. intricata* comb. nov. and *Hyd. rigidula* comb. nov.

Taxonomy

This section includes morphological description of two new species, *Hyd. orienticorrugata* sp. nov. and *Hyd. subtabacina* sp. nov. and a previously unreported species in Korea, *Hyd. rimosa*.


Hydnoporia subtabacina He390 CN "Hyd. tabacina"
Hydnoporia subtabacina Otto Miettinen 17028.3 (H) US-NY "Hyd. tabacina"
Hydnoporia subtabacina Viacheslav Spirin 6507 (H) RU-KHA "Hyd. tabacina"
Hydnoporia subtabacina CFMR:DLL2011-175 US-WI "Hyd. tabacina"
Hydnoporia subtabacina Heikki Kotiranta 20797 (H) RU-PER "Hyd. tabacina"
Hydnoporia subtabacina Viacheslav Spirin 6566 (H) RU-KHA "Hyd. tabacina"
Hydnoporia subtabacina SFC20190322-02 KR^T
Hydnoporia subtabacina SFC20190510-01 KR
Hydnoporia subtabacina Heikki Kotiranta 25205 (H) RU-KK "Hyd. tabacina"
Hydnoporia subtabacina Viacheslav Spirin 5196 (H) RU-KHA "Hyd. tabacina"
Hydnoporia subtabacina Viacheslav Spirin 6520 (H) RU-KHA "Hyd. tabacina"
Hydnoporia tabacina A. M. Ainsworth 2017-1-17 (K(M) 233332) GB
Hydnoporia tabacina Viacheslav Spirin 6066a (H) RU-NIZ
Hydnoporia tabacina Otto Miettinen 22126 (H) FI
Hydnoporia rhododendri N. Gerhold 2005-6-3 AT
Hydnoporia rhododendri Viacheslav Spirin 6476 (H) RU-PRI
Hydnoporia intricata He21064 CN
Hydnoporia intricata He1181 CN
Hydnoporia intricata KUC20121123-03 KR
Hydnoporia intricata KUC20211030-01 KR
Hydnoporia intricata KUC20210428-20 KR
Hydnoporia intricata SFC20160920-36 KR
Hydnoporia intricata SFC20170908-28 KR
Hydnoporia intricata SFC20140313-01 KR
Hydnoporia orienticorrugata KUC20131001-21 KR
Hydnoporia orienticorrugata He839 CN "Hyd. corrugata"
Hydnoporia orienticorrugata CLZhao938 CN-YN "Hyd. corrugata"
Hydnoporia orienticorrugata SFC20150212-01 KR
Hydnoporia orienticorrugata SFC20151030-12 KR^T
Hydnoporia yasudai KUC20180326-05 KR
Hydnoporia yasudai KUC20210319-14 KR
Hydnoporia yasudai SFC20160517-06 KR
Hydnoporia yasudai KUC20100409-18 KR
Hydnoporia yasudai SFC20150707-58 KR
Hydnoporia yasudai He273 CN
Hydnoporia yasudai CLZhao 1549 CN-YN
Hydnoporia yasudai Viacheslav Spirin 5533 (H) RU-KHA
Hydnoporia yasudai Viacheslav Spirin 6475 (H) RU-PRI
Hydnoporia yasudai SFC20160527-41 KR
Hydnoporia yasudai SFC20180410-24 KR
Hydnoporia yasudai SFC20180712-04 KR
Hydnoporia yasudai TNSF78711 JP
Hydnoporia yasudai SFC20160114-04 KR
Hydnoporia gigasetosa He1442 CN-YN
Hydnoporia gigasetosa He1461 CN-YN
Hydnoporia latesetosa He502 CN-HI^T
Hydnoporia latesetosa He492 CN-HI
Hydnoporia subrigidula He1157 CN-YN^T
Hydnoporia subrigidula He1123 CN-YN
Hydnoporia rigidula He379 CN
Hydnoporia rigidula SFC20140723-16 KR
Hydnoporia rigidula SFC20170324-10 KR
Hydnoporia lamellata Dai10527 CN
Hydnoporia lamellata Cui7629 CN
Hydnoporia tabacinoides Cui10428 CN
Hydnoporia tabacinoides CLZhao859 CN-YN
Hydnoporia rimosa Viacheslav Spirin 5277 (H) RU-KHA
Hydnoporia rimosa Viacheslav Spirin 6104 (H) RU-KHA
Hydnoporia rimosa KUC20121109-19 KR
Hydnoporia rimosa Viacheslav Spirin 5678 (H) RU-KHA
Hydnoporia diffissa Otto Miettinen 19463 (H 7008917) US-NY^T
Hydnoporia corrugata A3_wood_2 GB
Hydnoporia corrugata Jon Klepsland 2011-7-24 (O F-247869) NO^T
Hydnoporia diffissa Otto Miettinen 17127.4 (H) US-NC
Hydnoporia olivacea P1201B PE
Hydnoporia olivacea Otto Miettinen & Kelo Käppi 16044 (H 7005770) US-MA^T
Hydnoporia laricicola Viacheslav Spirin 5400 (H) RU-KHA
Hydnoporia laricicola Dai13458 CN-HL^T
Porodaedalea alpicola Cui 12280 CN
0.05
Hydnoporia subtabacina sp. nov.
Hydnoporia intricata comb. nov.
Hydnoporia orienticorrugata sp. nov.
Hydnoporia yasudai
Hydnoporia rigidula comb. nov.
Hydnoporia rimosa

Figure 2. BI tree based on the ITS and *tef1* concatenated sequence datasets. The node numbers indicate the posterior probabilities (PP) above 0.7. *Hydnoporia* specimens examined in this study are shown in bold. Newly generated sequences in this study are shown in blue and bold. *Porodaedalea alpicola* (Cui 12280) is used as an outgroup. Letter codes after specimen voucher indicate ISO 3166 country code followed by the origin province. Detailed information is in Table 1. Type specimens are indicated with "T".

Hydnoporia orienticorrugata **M.Cho, Y.Cho, Y.W.Lim & J.J.Kim, sp. nov.**

MycoBank No: 854671

Fig. 3

Diagnosis. Resupinate, effused basidiome, smooth, brown to reddish brown hymenial surface, sterile margin; cylindrical basidia with 10.3–15.9 × 2.8–4.0 µm, sharp-pointed setae with widened basal part and 35.0–64.6 × 8.1–13.2 µm, narrowly cylindrical to allantoid basidiospores with 4.4–5.8 × 1.5–2.0 µm, and growing on an angiosperm branch (a few on gymnosperm branches).

Type. Korea • Gangwon-do, Pyeongchang-gun, Mt. Heungjeong, 37°65.71'N, 128°32.25'E, alt. 800 m, 30 Oct 2015, Y. W. Lim, (***holotype*:** NIBRFG0000516804; ***isotype*:** SFC20151030-12).

Description. ***Basidiome*** resupinate, effused, thin, covering up to 0.1 mm thick. ***Hymenial surface*** smooth, membranaceous, crustaceous, with many cracks, brown (7.5YR, 4/2) to reddish brown (5YR, 5/3). ***Margin*** sterile, even, concolorous with that of the center. ***Hyphal system*** dimitic; generative hyphae septate, branched, without a clamp connection, thick-walled, few thin-walled, hyaline, 2.7–3.5 µm. Skeletal hyphae aseptate, unbranched, without a clamp connection, thick-walled, reddish-yellow (5YR, 7/8) to yellow (10YR, 7/8), 3.4–4.4 µm.



Figure 3. Morphological characteristics of *Hydnoporia orienticorrugata* (NIBRFG0000516804, holotype) **A** basidiome **B** microscopic features, **s** basidiospores; **b1** basidia; **b2** basidioles; **h1** generative hyphae; **h2** skeletal hyphae; **se** setae. Scale bars: 10 μm (**B**).

Basidia cylindrical, 4-spored, smooth, thin-walled, hyaline, 10.3–15.9(–16.6) × 2.8–4.0 μm. ***Setae*** sharp-pointed, few blunt-pointed, encrusted with crystals, distinctly widened basal part, smooth, bearing narrow lumen, thick-walled, reddish-brown (2.5YR, 4/4), 35.0–64.6 × 8.1–13.2(–14.2) μm. ***Basidiospores*** narrowly cylindrical to allantoid, smooth, slightly curved, thin-walled, hyaline, with narrow apex, a few bearing oil droplets, 4.4–5.8(–6.1) × 1.5–2.0 μm, L = 5.10 μm, W = 1.72 μm, Q = 2.53–3.44, n = 21.

Distribution. East Asia (Korea, China).

Ecology. Grew on an angiosperm branch in mixed hardwood forest, although a few grew on gymnosperm branches.

Etymology. Named after its distribution in East Asian regions and morphological similarity to *Hydnoporia corrugata*.

Additional specimens examined. Korea • Gangwon-do, Pyeongchang-gun, Odaesan National Park, 37°44.06'N, 128°35.25'E, alt. 690 m, 19 Oct 2012, Y. Jang & S. Jang, KUC20121019-16; Korea • Gangwon-do, Pyeongchang-gun, Odaesan National Park, 37°44.30'N, 128°35.03'E, alt. 660 m, 23 Nov 2012, Y. Jang & S. Jang, KUC20121123-05; Korea • Gangwon-do, Pyeongchang-gun, Odaesan National Park, 37°44.04'N, 128°35.03'E, alt. 680 m, 01 Oct 2013, Y. Jang & S. Jang, KUC20131001-21; Korea • Gangwon-do, Injae-gun, Mt. Bangtae, 37°87.53'N, 128°31.12'E, alt. 390 m, 12 Feb 2015, Y. W. Lim, SFC20150212-01.

Notes. Our specimens were phylogenetically well grouped with the Chinese specimens (He 761, He 839, and CLZhao 938), which were labeled either as *Hymenochaete corrugata*, *Hymenochaetopsis corrugata*, or *Pseudochaete corrugata* (He and Dai 2012; Yang et al. 2016) (Figs 1, 2). Other than these three specimens and KUC20121019-16 from Korea, no other records were found in East Asia, even when the other synonyms of *H. corrugata* were considered. Nevertheless, the East Asian clade formed a distinct clade from the clade with European *Hyd. corrugata*, which included the neotype specimen from Norway (O F-247869). Our findings conform to those of an earlier study, which suggested that sequences identified as *Hyd. corrugata* in Korea and China could represent a novel species (Miettinen et al. 2019). *Hydnoporia orienticorrugata* sp. nov. has micromorphological characteristics similar to those of *Hyd. corrugata*, but the latter has a grey to pale brown hymenial surface (Miettinen et al. 2019) that differs from the new species. Additionally, *Hyd. orienticorrugata* occurs on angiosperm and gymnosperm branches in Korea and China (He and Dai 2012) whereas *H. corrugata* occurs only on angiosperm branches and seems to be restricted to Europe (Austria, England, Ireland, Norway, Russia, and Sweden) (Fries 1815; Grundy et al. 2012; Miettinen et al. 2019).

Hydnoporia subtabacina **M.Cho, Y.Cho, Y.W.Lim & J.J.Kim, sp. nov.**

MycoBank No: 854672

Fig. 4

Diagnosis. Effused-reflexed, pileate basidiome, smooth, brown hymenial surface, sterile margin; cylindrical basidia with 14.6–17.9 × 2.9–3.8 µm, sharp-pointed and few elongated setae with 58.6–140.0 × 9.8–26.1 µm, narrowly cylindrical basidiospores with 4.4–5.7 × 1.6–1.9 µm, and occurs on angiosperm trees (branches and trunks).

Type. Korea • Gyeongsangbuk-do, Bonghwa-gun, Taebaeksan National Park, Baekcheon valley, 37°00.64'N, 128°98.41'E, alt. 830 m, 22 Mar 2019, Y. W. Lim & S. Yoo, (***holotype*****:** NIBRFG0000505378; ***isotype*****:** SFC20190322-02).

Description. ***Basidiome*** effused-reflexed, pileate, 0.1 mm thick. ***Hymenial surface*** smooth, membranaceous, brown (7.5YR, 5/4) to dark brown (7.5YR, 3/3). ***Margin*** sterile, slightly lighter. ***Hyphal system*** dimitic; generative hyphae septate, frequently branched, without a clamp connection, thick-walled, hyaline, 2.3–3.8 µm. Skeletal hyphae aseptate, unbranched, without a clamp connection, thick-walled, reddish-yellow (5YR, 7/8) to yellow (10YR, 7/8), 3.0–5.0 µm.

Basidia cylindrical, 4-spored, smooth, thin-walled, hyaline, (14.0–)14.6–17.9(–18.7) × 2.9–3.8 µm. ***Setae*** sharp-pointed, encrusted with crystals, cylindrical, fusiform, few elongated apex, smooth, bearing a narrow or wide lumen, thick-walled, dark reddish brown (5YR, 3/4) to dusky red (2.5YR, 3/2), 58.6–140.0 × 9.8–26.1 µm. ***Basidiospores*** narrowly cylindrical, smooth, slightly curved, thin-walled, hyaline, with narrow apex, 4.4–5.7 × 1.6–1.9 µm, L = 5.22 µm, W = 1.68 µm, Q = 2.78–3.61, n = 21.

Distribution. Korea, China, Far East Russia, and US.

Ecology. Grew on *Quercus* in angiosperm forest.



Figure 4. Morphological characteristics of *Hydnoporia subtabacina* (NIBRFG0000505378, **holotype**) **A** basidiome **B** microscopic features, **s** basidiospores; **b1** basidia; **b2** basidioles; **h1** generative hyphae; **h2** skeletal hyphae; **se** setae. Scale bars: 10 µm (**s**, **b1**, **b2**, **h1**, **h2**); 20 µm (**se**).

Etymology. Named after its morphological similarity with *Hydnoporia tabacina*.

Additional specimens examined. KOREA • Gangwon-do, Taebaek-si, Taebaeksan National Park, Yuilsa Temple, 37°10.87'N, 128°91.07'E, alt. 1,250 m, 10 May 2019, Y. W. Lim & S. Yoo, SFC20190510-01.

Notes. According to our phylogenetic analysis, sequences annotated as '*Hyd. tabacina*' were divided into a European and an Asian-North American clade (Fig. 2). The holotype specimen sequence is unavailable, but it is known that the specimen (≡*Aricularia tabacina* Sowerby) locality is Britain (Sowerby 1797) and the lectotype specimen is from Sweden (Miettinen et al. 2019). Therefore, we acknowledge the European clade (Finland, Great Britain, and Western Russia) as *Hyd. tabacina* and the Asian-American clade (China, Fareast Russia, Korea, and the US) as the new species, following the results of a previous study (Miettinen et al. 2019). *Hydnoporia subtabacina* sp. nov. occurs only on angiosperm branches or trunks. The microscopic characteristics of *Hyd. subtabacina* and *Hyd. tabacina* are similar, but longer and wider basidiospores are reported in the latter species, viz. 4.58–5.9 × 1.78–2.02 µm (Miettinen et al. 2019). Further, setal measurements of our specimens had broader variation (58.6–140.0 × 9.8–26.1 µm) compared to those of the Fareast Russian (63.92–94.15 × 9.38–14.5 µm) and North American specimens (71.1–97.9 × 9.5–14.23 µm) (Miettinen et al. 2019). While there is a morphological description of Chinese '*Hyd. tabacina*', no sequence data were available for these observed specimens

(Dai 2010). Additionally, the morphological characteristics of the Chinese '*Hyd. tabacina*' with a hydnoid and yellowish basidiome (Dai 2010) differ from those of the European specimens and of *Hyd. subtabacina*. Therefore, further research is needed for an accurate identification of Chinese '*Hyd. tabacina*'.

***Hydnoporia rimosa* (Lloyd) Spirin & Miettinen, Fungal Systematics and Evolution 4: 92 (2019)**

MycoBank No: 830597

Fig. 5

Diagnosis. Resupinate, effused basidiome, reddish brown hymenial surface, white to brown margin; cylindrical to narrowly clavate basidia with 10.0–14.1 × 2.7–3.4 µm, sharp to blunt pointed setae with 44.4–83.1 × 8.2–13.4 µm, narrowly cylindrical to allantoid basidiospores with 4.7–6.1 × 1.7–2.0 µm, and occurs on angiosperm branches.

Type. JAPAN • Tohoku, Sendai, 24 Oct 1920, Yasuda, (***holotype*:** TNS-F203210; ***lectotype*:** MBT387146).

Description. ***Basidiome*** resupinate, effused, thin, leathery, up to 0.2 mm thick. ***Hymenial surface*** smooth, membranaceous, crustaceous, with many cracks, light reddish-brown (2.5YR, 6/4) to reddish-brown (2.5YR, 5/3). ***Margin*** sterile, even, edge whitish (7.5YR, 9/2) when fresh and becomes brown (7.5YR, 5/8). ***Hyphal system*** dimitic; generative hyphae septate, branched, without a clamp connection, thick-walled, hyaline to pale brown (10YR, 6/3), 3.0–3.5 µm. Skeletal hyphae aseptate, unbranched, without a clamp connection, thick-walled, yellowish (7.5YR, 6/8) to reddish brown (2.5YR, 4/3), 3.5–4.2 µm.

Basidia cylindrical to narrowly clavate, 4-spored, smooth, thin-walled, hyaline, 10.0–14.1(–14.5) × 2.7–3.4 µm. ***Setae*** sharp- to blunt-pointed, encrusted with crystals, smooth, bearing a wide or narrow lumen, thick-walled, projecting up to 15 µm above the hymenium, dark reddish brown (2.5YR, 2.5/3), 44.4–83.1 × 8.2–13.4(–18.6) µm. ***Basidiospores*** narrowly cylindrical to allantoid, smooth, thin-walled, hyaline, with narrow apex, 4.7–6.1 × (1.5–)1.7–2.0 µm, L = 5.40 µm, W = 1.80 µm, Q = 2.64–3.30, n = 20.

Specimen examined. KOREA • Gangwon-do, Pyeongchang-gun, Odaesan National Park, 37°44.30'N, 128°35.01'E, alt. 660 m, 9 Nov 2012, Y. Jang & S. Jang, KUC20121109-19.

Notes. The observed *Hyd. rimosa* specimen from Korea has similar morphological characteristics as those of the type specimen (TNS-F203210), but the type specimen has abundant blunt-pointed setae and wider basidiospores (L = 5.22, W = 1.99, Q = 2.47–2.78) (Miettinen et al. 2019) compared to the Korean specimen. In the phylogenetic tree, *Hyd. diffissa* is closely related to *Hyd. rimosa*. These two species are practically indistinguishable in morphology, except that *Hyd. rimosa* has a wider lumen and greater size variation of setae (Miettinen et al. 2019). However, they are geographically distinct, where *Hyd. diffissa* is distributed across North and South America (Colombia, Peru, and Eastern US) (Miettinen et al. 2019), while *Hyd. rimosa* is distributed in East Asia (Japan, Korea, and Far East Russia) (Ito 1930b; Miettinen et al. 2019).



Figure 5. Morphological characteristics of *Hydnoporia rimosa* (KUC20121109-19) **A** basidiome **B** microscopic features, **s** basidiospores; **b1** basidia; **b2** basidioles; **h1** generative hyphae; **h2** skeletal hyphae; **se** setae. Scale bars: 10 µm (**B**).

***Hydnoporia intricata* (Lloyd) M.Cho, Y.Cho, Y.W.Lim & J.J.Kim, comb. nov.**

MycoBank No: 854673

Stereum intricatum Lloyd, Mycol. Writ. 7(67): 1157, 1922. Basionym.

Hymenochaete intricata (Lloyd) T.Ito, in Tokyo, Bot. Mag. 44: 156, 1930. Synonyms.

=*Pseudochaete intricata* (Lloyd) S.H.He & Y.C.Dai, Fungal Diversity, 56: 89, 2012.

=*Hymenochaetopsis intricata* (Lloyd) S.H.He & Jiao Yang, Mycol. Prog. 15(2/13): 6, 2016.

***Hydnoporia rigidula* (Berk. & M.A.Curtis) M. Cho, Y.Cho, Y.W.Lim & J.J.Kim, comb. nov.**

MycoBank No: 854674

Hymenochaete rigidula Berk. & M.A.Curtis, Journal of the Linnean Society. Botany 10: 334, 1869. Basionym.

Pseudochaete rigidula (Berk. & M.A.Curtis) S.H.He & Y.C.Dai, Fungal Diversity 56: 89, 2012. Synonyms.

=*Hymenochaetopsis rigidula* (Berk. & M.A.Curtis) S.H.He & Jiao Yang, Mycol. Prog. 15 (2/13): 6, 2016.

Taxonomic key to *Hydnoporia* in Korea

1 Basidiome resupinate **2**
– Basidiome effused-reflexed **5**
2 Basidiome margin strictly attached to substrate **3**
– Basidiome margin detached from substrate **4**
3 Setae sharp-pointed, distinctly widened base ..... ***Hyd. orienticorrugata***
– Setae sharp– to blunt-pointed, narrow base ..... ***Hyd. rimosa***
4 Occurs on dead angiosperm branches, setae 29.1–66.4 × 5.0–9.2 µm ..... ***Hyd. rigidula***
– Occurs on dead gymnosperm branches, setae 36.0–92.7 × 9.1–19.6 µm ..... ***Hyd. yasudai***
5 Basidiospores cylindrical, Q value < 3.0 ..... ***Hyd. subtabacina***
– Basidiospores allantoid, Q value > 3.0 ..... ***Hyd. intricata***

Discussion

Some *Hydnoporia* species are indistinguishable based on a phylogenetic tree that is inferred from ITS data alone, notably as *Hyd. rhododendri*, *Hyd. tabacina*, and *Hyd. subtabacina* (Fig. 1). This is resolved by including an additional genetic marker, *tef1*, to infer a multigenetic marker phylogeny (Fig. 2). A phylogenetic analysis solely based on ITS for fungal species identification has been criticized by taxonomists (Harder et al. 2013; Santos et al. 2017). The use of ITS alone works well for several genera (Frøslev et al. 2007; Hallenberg et al. 2007). However, it may lead to under-splitting of some taxa (Harder et al. 2013) or over-splitting of other, taxa as seen for *Hyd. yasudai* (Fig. 1). Therefore, the use of additional protein-coding genetic markers for phylogenetic analyses is essential to achieve properly resolved species clades (Fig. 2, Suppl. material 1).

To the 14 previously accepted species in *Hydnoporia*, the present study adds four ones: two new *Hydnoporia* species were described, and two other species were transferred to *Hydnoporia*. Of these 18 species, 13 have been reported in East Asia (China, Japan, Korea, and Fareast Russia), namely *Hydnoporia gigasetosa*, *Hyd. intricata*, *Hyd. lamellata*, *Hyd. laricicola*, *Hyd. latesetosa*, *Hyd. orienticorrugata* sp. nov., *Hyd. rhododendri*, *Hyd. rigidula*, *Hyd. rimosa*, *Hyd. subrigidula*, *Hyd. subtabacina* sp. nov., *Hyd. tabacinoides*, and *Hyd. yasudai* (Miettinen et al. 2019). Regarding the remaining five species, *Hyd. diffissa*, *Hyd. lenta*, and *Hyd. olivacea* have only been reported in the Americas, and *Hyd. corrugata* and *Hyd. tabacina* have only been reported in Europe (Miettinen et al. 2019). This indicates that many *Hydnoporia* species are geographically or ecologically restricted unlike many other wood-decaying fungi that are cosmopolitan and less constrained by environmental factors (Sato et al. 2012). The regional constraint for *Hydnoporia* may be the consequence of host or vector specificity.

In Korea, three previously recorded species – namely *Hydnoporia intricata*, *Hyd. rigidula*, and *Hyd. yasudai* – have been phylogenetically verified. Clades of the first two species were supported by high bootstrap values and posterior probabilities. However, *Hyd. yasudai* formed a complex, as observed by Miettinen et al. (2019), who suggested dividing the *Hyd. yasudai* complex into three to six different species. In our study, *Hyd. yasudai* specimens had large sequence variations in ITS (13 bp, 1.3%) and *tef1* (20 bp, 3.4%) but morphologically, the spore sizes

were relatively constant among the Fareast Russian, Japanese, and Korean specimens (Miettinen et al. 2019). Additionally, *Hyd. yasudai* has a specific host preference for gymnosperms (*Pinus* spp., *Abies firma*, and *A. holophylla*, etc.) (Léger 1998; Dai 2010; Miettinen et al. 2019). Therefore, we proposed the *Hyd. yasudai* complex to remain as a single species, with variation based on geographical distribution. Two *Hydnoporia* species previously reported in Korea, *Hyd. corrugata* and *Hyd. tabacina*, were each represent other species. East Asian '*Hyd. corrugata*' specimens (Korean and Chinese) (He and Dai 2012) were phylogenetically separated from the European *Hyd. corrugata*, which contains the type specimen (O F-247869) (Figs 1, 2), and it was thus introduced as a new species, *Hyd. orienti-corrugata* sp. nov. This result was consistent with that of Miettinen et al. (2019). Similarly, East Asian and North American '*Hyd. tabacina*' were different from the holotype – a European specimen – both phylogenetically and morphologically (Figs 2, 4), and they were introduced as a new species, *Hyd. subtabacina* sp. nov.

Two species combinations are proposed, viz. *Hyd. intricata* comb. nov. and *Hyd. rigidula* comb. nov. *Hydnoporia intricata* was first described as *Stereum intricatum* by Lloyd in 1922. It was then renamed to *Hymenochaete intricata* (Ito 1930a). After decades, He & Dai renamed it to *Pseudochaete intricata* (He and Dai 2012) and then Yang suggested *Hymenochaetopsis intricata* (Yang et al. 2016). Based on a recent study, *Pseudochaete* and *Hymenochaetopsis* are no longer valid and are considered younger synonyms of *Hydnoporia* (Miettinen et al. 2019). Based on this study, *Hymenochaete intricata* is verified to belong to *Hydnoporia*. The morphological characteristics of the Korean *Hyd. intricata* specimens studied here correspond to those of the original description. Therefore, we suggest that *Hymenochaete intricata* should be included in *Hydnoporia*. This result was further supported by phylogenetic analysis of combined sequence datasets (ITS+*tef1*) with high bootstrap support value and posterior probability (Fig. 2, Suppl. material 1).

Hydnoporia rigidula was initially reported as *Hymenochaete rigidula* Berk. & M. A. Curtis in 1868 (Berkeley and Curtis 1868). Based on multiple taxonomic revisions, it was renamed *Pseudochaete rigidula* (He and Dai 2012) and later *Hymenochaetopsis rigidula* according to (Yang et al. 2016). According to Miettinen et al. (2019) and this study, *Hymenochaete rigidula* is phylogenetically located in *Hydnoporia*. Miettinen et al. (2019) also stated that East Asian *Hyd. rigidula* may be distinct from the American *Hyd. rigidula*. However, no sequenced specimens are available from the American or Caribbean (Cuba and Jamaica) regions (He and Dai 2012; Yang et al. 2016) to verify the differences. Nevertheless, East Asian and American specimens have similar setae and basidiospore size measurements and other macro-morphological characteristics (Parmasto 2001; He and Dai 2012; Kim et al. 2016). Therefore, further assessment is required to separate the species.

In conclusion, we propose two new species and two species combinations within the genus *Hydnoporia*. Given the morphological similarities between *Hydnoporia* and *Hymenochaete*, molecular analysis is crucial for accurate species identification, ideally using multiple genetic regions. This study resolved the taxonomic confusion arising from the continuous systematic revision of some *Hydnoporia* species and emphasized the need to update old names to avoid confusion. Although *Hydnoporia* appears to be primarily distributed in East Asia, with a few species in Europe and the Americas, it remains underexplored in the Southern Hemisphere. Therefore, further investigation of the

global distribution and biogeographical relationships of *Hydnoporia* is necessary to understand the true diversity of the genus and establish a stable species classification.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was supported by the project of the National Institute of Biological Resources [NIBR202402104] under the Ministry of Environment, Republic of Korea. This study was also supported by the Korea University Grant.

Author contributions

Conceptualization: MC, YC, YWL. Data curation/collection: MC, YC, DK. Investigation: KH, YWL, JJK. Methodology: MC, YC, SLK. Project administration: YWL, JJK. Funding acquisition: JJK. Writing – original draft: MC, YC. Writing – review & editing: MC, YC, SLK, DK, YWL, JJK.

Author ORCIDs

Minseo Cho https://orcid.org/0009-0007-0834-7294
Yoonhee Cho https://orcid.org/0000-0002-0743-0914
Sun Lul Kwon https://orcid.org/0000-0002-2075-5342
Dohye Kim https://orcid.org/0009-0007-0551-099X
Kentaro Hosaka https://orcid.org/0000-0002-4469-8303
Young Woon Lim https://orcid.org/0000-0003-2864-3449
Jae-Jin Kim https://orcid.org/0000-0001-8990-2139

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

Berkeley MJ, Curtis MA (1868) Fungi Cubenses (Hymenomycetes). Journal of the Linnean Society of London, Botany 10(45): 280–320. https://doi.org/10.1111/j.1095-8339.1868.tb00529.x

Brazee NJ, Lindner DL, D'Amato AW, Fraver S, Forrester JA, Mladenoff DJ (2014) Disturbance and diversity of wood-inhabiting fungi: Effects of canopy gaps and downed woody debris. Biodiversity and Conservation 23(9): 2155–2172. https://doi.org/10.1007/s10531-014-0710-x

Color BM (2009) Munsell soil-color charts with genuine Munsell® color chips. Munsell Color, Grand Rapids (MI).

Connell L, Staudigel H (2013) Fungal diversity in a dark oligotrophic volcanic ecosystem (DOVE) on Mount Erebus, Antarctica. Biology 2(2): 798–809. https://doi.org/10.3390/biology2020798

Dai Y-C (2010) Hymenochaetaceae (Basidiomycota) in China. Fungal Diversity 45(1): 131–343. https://doi.org/10.1007/s13225-010-0066-9

Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: More models, new heuristics and high-performance computing. Nature Methods 9(8): 772. https://doi.org/10.1038/nmeth.2109

Fries EM (1815) Observationes mycologicae. G. Bonnieri, Copenhagen, Denmark. https://doi.org/10.5962/bhl.title.112534

Frøslev TG, Jeppesen TS, Læssøe T, Kjøller R (2007) Molecular phylogenetics and delimitation of species in *Cortinarius* section *Calochroi* (Basidiomycota, Agaricales) in Europe. Molecular Phylogenetics and Evolution 44(1): 217–227. https://doi.org/10.1016/j.ympev.2006.11.013

Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. Molecular Ecology 2(2): 113–118. https://doi.org/10.1111/j.1365-294X.1993.tb00005.x

Grundy KC, Woodward S, Genney DR, Taylor AF (2012) A molecular approach to explore the extent of the threatened fungus *Hypocreopsis rhododendri* within wood. Fungal Biology 116(3): 354–362. https://doi.org/10.1016/j.funbio.2011.12.002

Hallenberg N, Nilsson RH, Antonelli A, Wu SH, Maekawa N, Nordén B (2007) The *Peniophorella praetermissa* species complex (Basidiomycota). Mycological Research 111(12): 1366–1376. https://doi.org/10.1016/j.mycres.2007.10.001

Harder CB, Læssøe T, Frøslev TG, Ekelund F, Rosendahl S, Kjøller R (2013) A three-gene phylogeny of the *Mycena pura* complex reveals 11 phylogenetic species and shows ITS to be unreliable for species identification. Fungal Biology 117(11–12): 764–775. https://doi.org/10.1016/j.funbio.2013.09.004

He S-H, Dai Y-C (2012) Taxonomy and phylogeny of *Hymenochaete* and allied genera of Hymenochaetaceae (Basidiomycota) in China. Fungal Diversity 56(1): 77–93. https://doi.org/10.1007/s13225-012-0174-9

He S-H, Li H-J (2013) *Pseudochaete latesetosa* and *P. subrigidula* spp. nov. (*Hymenochaetales*, Basidiomycota) from China based on morphological and molecular characters. Mycological Progress 12(2): 331–339. https://doi.org/10.1007/s11557-012-0838-6

Ito T (1930a) Symbolæ ad Mycologiam Japonicam. V. *Hymenochaete*. Botanical Magazine Tokyo 44(519): 151–157. https://doi.org/10.15281/jplantres1887.44.151

Ito T (1930b) Symbolæ ad Mycologiam Japonicam. IV. *Asterostromella* et *Hymenochaete*. Botanical Magazine Tokyo 44(518): 89–93. https://doi.org/10.15281/jplantres1887.44.89

Jang Y, Jang S, Min M, Hong J-H, Lee H, Lee H, Lim YW, Kim J-J (2015) Comparison of the diversity of basidiomycetes from dead wood of the Manchurian fir (*Abies holophylla*) as evaluated by fruiting body collection, mycelial isolation, and 454 sequencing. Microbial Ecology 70(3): 634–645. https://doi.org/10.1007/s00248-015-0616-5

Jang Y, Jang S, Lee J, Lee H, Lim YW, Kim C, Kim J-J (2016) Diversity of wood-inhabiting polyporoid and corticioid fungi in Odaesan National Park, Korea. Mycobiology 44(4): 217–236. https://doi.org/10.5941/MYCO.2016.44.4.217

Jeong WJ, Lim YW, Lee JS, Jung HS (2005) Phylogeny of *Phellinus* and related genera inferred from combined data of ITS and mitochondrial SSU rDNA sequences. Journal of Microbiology and Biotechnology 15: 1028–1038.

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. https://doi.org/10.1093/molbev/mst010

Kim NK, Park JY, Park MS, Lee H, Cho HJ, Eimes JA, Kim C, Lim YW (2016) Five new wood decay fungi (*Polyporales* and *Hymenochaetales*) in Korea. Mycobiology 44(3): 146–154. https://doi.org/10.5941/MYCO.2016.44.3.146

Léger J-C (1998) Le genre *Hymenochaete* Léveillé. Bibliotheca Mycologica, 171. [J. Cramer: Berlin, Stuttgart, Germany.]

Miettinen O, Larsson K-H, Spirin V (2019) *Hydnoporia*, an older name for *Pseudochaete* and *Hymenochaetopsis*, and typification of the genus *Hymenochaete* (*Hymenochaetales*, Basidiomycota). Fungal Systematics and Evolution 4: 77–96. https://doi.org/10.3114/fuse.2019.04.07

Miller MA, Schwartz T, Pickett BE, He S, Klem EB, Scheuermann RH, Passarotti M, Kaufman S, O'Leary MA (2015) A RESTful API for access to phylogenetic tools via the CIPRES Science Gateway. Evolutionary Bioinformatics Online 11: 43–48. https://doi.org/10.4137/EBO.S21501

Noris F, Siegel JA, Kinney KA (2011) Evaluation of HVAC filters as a sampling mechanism for indoor microbial communities. Atmospheric Environment 45(2): 338–346. https://doi.org/10.1016/j.atmosenv.2010.10.017

Palla B, Yuan Y, Dai Y-C, Papp V (2023) Host preferences of *Pinus*-dwelling Hymenochaetaceae. In: Sridhar KR, Deshmukh SK (Eds) Ecology of Macrofungi: An Overview. CRC Press, 14 pp. https://doi.org/10.1201/9781003429272-13

Parmasto E (1995) The genus *Hymenochaete* (Hymenomycetes): Infrageneric classification and satellite genera. Documents Mycologiques 25: 305–315.

Parmasto E (2001) Hymenochaetoid fungi (Basidiomycota) of North America. Mycotaxon 79: 107–176.

Parmasto E, Saar I, Larsson E, Rummo S (2014) Phylogenetic taxonomy of *Hymenochaete* and related genera (Hymenochaetales). Mycological Progress 13(1): 55–64. https://doi.org/10.1007/s11557-013-0891-9

Rambaut A (2018) FigTree, a graphical viewer of phylogenetic trees (Version 1.4.4). Institute of evolutionary biology, University of Edinburgh, Edinburgh, UK.

Rehner SA, Buckley E (2005) A *Beauveria* phylogeny inferred from nuclear ITS and EF1–α sequences: Evidence for cryptic diversification and links to *Cordyceps* teleomorphs. Mycologia 97(1): 84–98. https://doi.org/10.3852/mycologia.97.1.84

Rittenour WR, Ciaccio CE, Barnes CS, Kashon ML, Lemons AR, Beezhold DH, Green BJ (2014) Internal transcribed spacer rRNA gene sequencing analysis of fungal diversity in Kansas City indoor environments. Environmental Science. Processes & Impacts 16(1): 33–43. https://doi.org/10.1039/C3EM00441D

Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19(12): 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Santos L, Alves A, Alves R (2017) Evaluating multi-locus phylogenies for species boundaries determination in the genus *Diaporthe*. PeerJ 5: e3120. https://doi.org/10.7717/peerj.3120

Sato H, Tsujino R, Kurita K, Yokoyama K, Agata K (2012) Modelling the global distribution of fungal species: New insights into microbial cosmopolitanism. Molecular Ecology 21(22): 5599–5612. https://doi.org/10.1111/mec.12053

Sayers EW, Cavanaugh M, Clark K, Pruitt KD, Sherry ST, Yankie L, Karsch-Mizrachi I (2024) GenBank 2024 update. Nucleic Acids Research 52(D1): D134–D137. https://doi.org/10.1093/nar/gkad903

Smith SA, Tank DC, Boulanger L-A, Bascom-Slack CA, Eisenman K, Kingery D, Babbs B, Fenn K, Greene JS, Hann BD, Keehner J, Kelley-Swift EG, Kembaiyan V, Lee SJ,

Li P, Light DY, Lin EH, Ma C, Moore E, Schorn MA, Vekhter D, Nunez PV, Strobel GA, Donoghue MJ, Strobel SA (2008) Bioactive endophytes warrant intensified exploration and conservation. PLoS ONE 3(8): e3052. https://doi.org/10.1371/journal.pone.0003052

Sowerby J (1797) Coloured Figures of English Fungi. Vol. 1, 54 pp. https://doi.org/10.5962/bhl.title.6342

Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30(9): 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Vu D, Groenewald M, De Vries M, Gehrmann T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. Studies in Mycology 92(1): 135–154. https://doi.org/10.1016/j.simyco.2018.05.001

Wagner T, Fischer M (2002) Classification and phylogenetic relationships of *Hymenochaete* and allied genera of the *Hymenochaetales*, inferred from rDNA sequence data and nuclear behaviour of vegetative mycelium. Mycological Progress 1(1): 93–104. https://doi.org/10.1007/s11557-006-0008-9

White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18: 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wu F, Dai S-J, Vlasák J, Spirin V, Dai Y-C (2019) Phylogeny and global diversity of *Porodaedalea*, a genus of gymnosperm pathogens in the *Hymenochaetales*. Mycologia 111(1): 40–53. https://doi.org/10.1080/00275514.2018.1526618

Wu F, Zhou L-W, Vlasák J, Dai Y-C (2022) Global diversity and systematics of Hymenochaetaceae with poroid hymenophore. Fungal Diversity 113(1): 1–192. https://doi.org/10.1007/s13225-021-00496-4

Yang J, Dai L-D, He S-H (2016) *Hymenochaetopsis* nom. nov. proposed to replace *Pseudochaete* (*Hymenochaetales*, Basidiomycota) with descriptions of *H. laricicola* sp. nov and *H. gigasetosa* new to China. Mycological Progress 15(2): 1–8. https://doi.org/10.1007/s11557-015-1153-9

Supplementary material 1

Data 1

Authors: Minseo Cho, Yoonhee Cho, Sun Lul Kwon, Dohye Kim, Kentaro Hosaka, Young Woon Lim, Jae-Jin Kim

Data type: fasta

Explanation note: ITS sequence alignments of all studied sequence datasets.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mycokeys.111.137347.suppl1

Supplementary material 2

Data 2

Authors: Minseo Cho, Yoonhee Cho, Sun Lul Kwon, Dohye Kim, Kentaro Hosaka, Young Woon Lim, Jae-Jin Kim

Data type: fasta

Explanation note: Concatenated multimarker (ITS+*tef1*) sequence alignments of studied sequence datasets.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mycokeys.111.137347.suppl2

Supplementary material 3

ML tree based on the ITS and *tef1* concatenated sequence datasets

Authors: Minseo Cho, Yoonhee Cho, Sun Lul Kwon, Dohye Kim, Kentaro Hosaka, Young Woon Lim, Jae-Jin Kim

Data type: png

Explanation note: The node numbers indicate the bootstrap support values (BS) above 70%. *Hydnoporia* specimens examined in this study are shown in bold. Newly generated sequences in this study are shown in blue and bold. *Porodaedalea alpicola* (Cui 12280) is used as an outgroup. Letter codes after specimen voucher indicate ISO 3166 country code followed by the province of origin. Type specimens are indicated with "T".

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/mycokeys.111.137347.suppl3